

Pre-Surgical Instruction: Medications To Avoid

DO NOT ingest any of the following for two week prior to surgery:

MEDICATIONS	HERBAL SUPPLEMENTS
<ul style="list-style-type: none">▪ 4-Way Cold Tablets▪ Advil Tablets and Capsules▪ Alka-Seltzer Effervescent Tablets▪ Alka-Seltzer Plus Cold Medicine Tablets▪ Aleve Tablets and Capsules▪ Anacin Tablets & Capsules, Any Strength▪ Aspergum▪ Arthritis Strength Bufferin▪ Bayer Aspirin Tablets▪ Bufferin Tablets▪ Bufferinol Tablets▪ Carisoprodol▪ Darvon▪ Doan's Pills▪ Ecotrin Tablets▪ Excedrin Tablets▪ Fiorinal Tablets▪ Ibuprofen Tablets & Capsules▪ Medipren Tablets & Capsules▪ Midol Capsules▪ Motrin Tablets & Capsules▪ Nuprin Tablets & Capsules▪ Pepto-Bismol Tablets & Suspension▪ Percodan & Percodan-Demi Tablets▪ St. Josephs Aspirin for Children▪ St. Josephs Cold Tablets for Children▪ Naproxen Tablets & Capsules▪ Soma▪ Talwin Compound	<ul style="list-style-type: none">▪ St. John's Wort▪ Ginseng▪ MaHuang▪ Kava Kava▪ Dong Quai▪ Cola Nut▪ Chastefruit tree▪ Yohimbe▪ Hawthorne▪ Eleurhero▪ Siberian Ginseng▪ All Energy Drinks